

CALORIMETRIC STUDY OF CHANGES INDUCED BY PRESERVATIVES IN LIQUID EGG PRODUCTS

Csaba Németh, Kinga Horváth, Árpád Drobecz, László Friedrich, Klára Pásztor-Huszár, Csaba Balla

Department of Refrigeration and Livestock Products Technology, Faculty of Food Science, Corvinus University of Budapest, Hungary

Key words: calorimetry, liquid egg, preservative, potassium sorbate, sodium benzoate

In our tests we aimed to study the effects of additives normally used in liquid egg products on the calorimetric properties of liquid egg samples. Raw liquid egg white ($\text{pH}=8.9\pm 0.1$), liquid egg yolk ($\text{pH}=6.5\pm 0.2$) and whole liquid egg ($\text{pH}=7.1\pm 0.2$) broken up in industrial conditions were used in our measurements. Sodium benzoate and potassium sorbate were added to the samples to achieve solutions of 0.1, 0.3 and 0.5 g/L concentrations. We used citric acid to prepare liquid egg products with various pH value (5.5, 5.0 and 4.5). The calorimetric tests were performed with MicroDSC III device. Liquid egg samples were heated up from 20°C to 95°C generally with a heating rate of 1.5°C/min. Sample weights were 500 mg \pm 0.1 mg and water was used as reference solution. Evaluation was performed by Seftsoft2000 software, a component of the device.

Our measurements have shown significant changes in calorimetric parameters of liquid egg samples by decreasing the pH value to 5.0. In addition to the reduced enthalpy due to acidification of samples, a decrease in denaturation temperature was also observed in the egg white. While the native egg white started to precipitate at 60°C, at pH 5.0 the denaturation was already started at 54.5°C. When sodium benzoate and potassium sorbate were added to liquid egg products a significant change of the tested calorimetric values was only found in the liquid egg samples containing the preservative at the concentration of 0.5 g/L.

INTRODUCTION

Many additives are used in the food industry. Some of them are added to the food for increased food safety to improve the efficiency of anti-microbial treatments or to extend shelf-life [Ogihova *et al.*, 2009; Rangan & Barceloux, 2009; Lee *et al.*, 2009]. Liquid egg manufacturers mainly use additives to extend shelf-life of these products. This is partly because customers using large quantities of liquid egg products prefer ready-for-use liquid egg products for the technology compared to powdered egg products, applicability and protein composition of which might be impaired [Landfeld *et al.*, 2008] but liquid egg products spoil rapidly even when stored in refrigerator.

When using additives, one should consider that calorimetric properties as well as heat sensitivity of liquid egg products (liquid whole egg, liquid egg white, and liquid egg yolk) may change [Torregiani *et al.*, 2009; Carmona *et al.*, 2007; Ichikawa & Schimomura, 2007; Mizutani *et al.*, 2006]. This can be a problem mainly in case of preservation technologies where the preservatives are added to the liquid egg prior to heat treatment. Such procedures include treatment of liquid eggs in packaging for avoidance of the chance of possible post-infection during the packaging process.

From egg components, proteins are the most heat sensitive; proteins are found in high percentage in the egg white [Hammershøj *et al.*, 2007; Rossi & Schiraldi, 1992; Gossett *et al.*, 1984]. Egg white is a protein system comprising ovomucin fibres incorporated into an aqueous solution containing

numerous globular proteins. The most important representatives of these proteins – due to their amount – include ovalbumin, conalbumin (ovotransferrin), ovomucoid, ovomucin, lysozyme and globulins [Chang *et al.*, 1977].

Composition of egg yolk is more complex; it can be best described as a complex system in which diverse particles are suspended in a protein solution (livetin) [Nielsen, 1998].

DSC method has already been used several times to study thermal denaturation of egg white and its fractions [Ferreira *et al.*, 1997; Donovan & Mapes, 1976; Donovan *et al.*, 1975]. Particularly, it helps measuring the enthalpy of denaturation (ΔH_d) in case of egg white and some of its components, and thereby it is able to provide quantitative information. Furthermore, denaturation temperature of various proteins can also be measured with this method [Andrássy *et al.*, 2006; Zhang *et al.*, 2004; Mohácsi-Farkas *et al.*, 1999].

In this study we investigated the calorimetric effects of potassium sorbate and sodium benzoate used in egg processing and confectionery industry [Gliemmo *et al.*, 2004] and pH-reduction with citric acid in raw liquid egg products homogenized in the routine industrial manner.

MATERIALS AND METHODS

Materials

Raw liquid egg white ($\text{pH}=8.9\pm 0.1$), egg yolk ($\text{pH}=6.5\pm 0.2$) and whole egg ($\text{pH}=7.1\pm 0.2$) broken under industrial conditions and homogenized in a piston-gap ho-

mogenizer at 100 bars were used in our tests. Sodium benzoate and potassium sorbate were added to the samples to achieve solutions of 0.1, 0.3, and 0.5 g/L concentrations. We used citric acid to prepare liquid egg products with various pH (5.5, 5.0, and 4.5).

CONSORT C831 model liquid pH-meter was used to control pH.

Differential Scanning Calorimetry (DSC)

Tests have been performed by using Setaram MicroDSC III device. Liquid egg samples were heated up from 20°C to 95°C at a heating rate of 1.5°C/min.

The measured mass of liquid egg samples was 500 mg±0.1mg and water was used as reference. In some cases a second test cycle was also performed but no reversible phenomenon was observed.

Evaluation was performed by Seftsoft2000 program, a component of the device.

RESULTS AND DISCUSSION

Analysis of homogenized egg products without preservative

First tests were carried out with raw liquid eggs without additives. Endothermic phenomena observed in the liquid egg white typically correspond to protein denaturation. However, from the 4 main components of egg white (conalbumin, lysozyme, ovalbumin, and globulins) fusion of peaks of lysozyme and conalbumin could be observed as a result of homogenization. Onset of denaturation was observed at 60°C while the peak (maximum) denaturation temperature was around 63.5°C (Figure 1).

The second peak (onset of denaturation at 70°C) is typical to denaturation of ovalbumin and globulins. Ratio of ovalbumin in liquid egg white is significantly higher (54% of total protein) compared to that of globulins (8% of total protein); therefore, peak parameters are determined by the calorimetric properties of ovalbumin. Peak denaturation temperature of these proteins was around 77°C (Figure 2).

Liquid egg yolk contains large amounts of lipoproteins. These cannot be separated well by calorimetric methods compared to the proteins of liquid egg white even in the non-homogenized sample. Therefore we analyzed one peak for each sample, *i.e.* total lipoproteins.




FIGURE 1. Thermogram of native liquid egg white without preservative (1. peak).




FIGURE 2. Thermogram of native liquid egg white without preservative (2. peak).

According to industrial experiences egg yolk is much less sensitive to heat than liquid egg white. Our tests have shown that peak denaturation temperature of these proteins was around 78°C (Figure 3).

The calorimetric properties of liquid whole egg (Figure 4) are influenced by the heat sensitive components of liquid egg white and egg yolk. Comparing Figures 2 and 3 it is apparent that peak denaturation temperatures of ovalbumin in the homogenized liquid egg white and liquid egg yolk lipoproteins was similar (77-78°C), and the thermogram of liquid whole egg is predominantly influenced by these two fractions. In the tests, we only investigated changes affecting significant technological applicability; therefore, in case of liquid whole egg we investigated only that peak.




FIGURE 3. Thermogram of native liquid egg yolk without preservative.




FIGURE 4. Calorimetric analysis of DSC curve of native liquid whole egg without preservative.

Analysis of homogenized egg products with preservative Baseline and peak denaturation point as well as enthalpy of various native liquid egg white fractions have changed by pH reduction with citric acid (Figure 5).

When pH of liquid egg white was reduced to 4.5 baseline denaturation point of the conalbumin-containing fraction reduced from 60°C to 45°C while denaturation enthalpy reduced by two-third (from 0.273 J/g to 0.092 J/g).

Denaturation temperatures of the ovalbumin-containing fraction considerably shifted. Reduction of pH of native liquid egg white (pH 8.0) to 4.5 reduced the baseline denaturation temperature from 72°C to 64°C and the peak denaturation temperature from 77°C to around 70°C. Nevertheless, denaturation enthalpy decreased only slightly by pH reduction (from 1.075 J/g to 0.965 J/g at pH 4.5).

In case of pH reduction of native liquid egg yolk (pH 7.0) with citric acid, a decrease in denaturation temperatures and denaturation enthalpy was observed (Figure 6).

For liquid egg yolk the baseline denaturation temperature has decreased to 64.5°C even at pH 5.5 compared to 69°C measured in native liquid egg yolk, but subsequently it did not change significantly. Contrarily, constant reduction in peak denaturation point has been found. While in native liquid egg white this value was around 78°C at pH 5.5, and reduced to around 74°C at pH 5.0, and 71°C at pH 4.5.




FIGURE 5. Changes in the thermogram of liquid egg white by pH modification.




FIGURE 6. Changes in the thermogram of liquid egg yolk by pH modification.

Changes in denaturation temperatures and denaturation enthalpy have been observed also in tests with liquid whole egg at various pH values (Figure 7). Baseline denaturation temperature has reduced from 70.5°C in the native liquid whole egg (pH 7.6) to below 65°C at pH 4.5 while peak denaturation point changed only to a slight extent (measured range 75-78°C).

Denaturation enthalpy decreased from 1.738 J/g to 0.975 J/g by pH reduction to 4.5. The most significant change in enthalpy was found with pH reduction from 5.5 to 5.0 resulting in enthalpy change from 1.704 g/J to 1.300 g/J.

Baseline and peak denaturation temperatures and enthalpy of egg products of reduced pH are summarized in Table 1, and 95% limits of confidence intervals are summarized in Table 3.

Assessment of calorimetric diagrams of native liquid eggs has shown results similar to literature data [Andrássy *et al.*, 2006; Ferreira *et al.*, 1997]. From the egg proteins characteristics of conalbumin were changed the most significantly. Cunningham & Lineweaver [1965] have also drawn this conclusion when they




FIGURE 7. Changes in the thermogram of liquid whole egg by pH modification.

TABLE 1. Changes in calorimetric properties of liquid egg product by pH reduction.

pH of liquid egg products	Liquid egg white		Liquid egg yolk	Liquid whole egg	
	Peak 1	Peak 2			
Native liquid egg	T(baseline)	60.34	71.93	69.14	70.49
	T(max)	63.52	77.06	78.35	77.91
	ΔH_d	0.2727	1.0750	1.7248	1.7384
pH 5.5	T(baseline)	60.00	70.58	64.66	68.01
	T(max)	62.47	76.46	77.75	77.94
	ΔH_d	0.1505	1.0598	1.4466	1.7040
pH 5.0	T(baseline)	54.47	67.77	64.42	67.04
	T(max)	57.31	73.73	73.90	75.13
	ΔH_d	0.1368	1.1155	1.1019	1.2229
pH 4.5	T(baseline)	45.12	63.72	64.25	64.78
	T(max)	48.99	70.28	70.79	76.76
	ΔH_d	0.0923	0.9647	0.4403	0.9753


T(baseline) – baseline denaturation temperature (°C); T(max) – peak denaturation temperature (°C); ΔH_d – change in denaturation enthalpy (J/g).

established that 40% of conalbumin is precipitated with heat treatment at 57°C for 10 min in a phosphate-bicarbonate buffer at pH 6. However, when conalbumin solution was adjusted to pH 9 and heated under the same conditions, the protein did not precipitate [Cunningham & Lineweaver, 1965].

When sodium benzoate and potassium sorbate were added to liquid egg no significant changes have been observed in calorimetric values compared to pH reduction; therefore, we only show DSC curves of liquid egg white samples that are more significant regarding the change in heat sensitivity (Figures 8 and 9).


No significant differences were measured in denaturation temperature for liquid egg white and egg yolk with preservative concentrations of 0.1 and 0.3 g/L. However, it is seen in Figure 9 that the course of the curve changed and it had two peaks meanwhile the position of denaturation peak was almost unchanged and the change in the area determining denaturation enthalpy was not significant either.

For liquid egg samples, the tests have shown smaller differences in denaturation temperature and enthalpy change of liquid egg products with added potassium sorbate and sodium benzoate compared to those with reduced pH (Tables 2 and 3), but we observed significant difference in liquid egg samples with 0.5 g/L



— 0.0 g/L potassium sorbate; - - - 0.1 g/L potassium sorbate; - · - 0.3 g/L potassium sorbate; ····· 0.5 g/L potassium sorbate

FIGURE 8. Changes in the thermogram of liquid egg white by addition of potassium sorbate.



— 0.0 g/L sodium benzoate; - - - 0.1 g/L sodium benzoate; - · - 0.3 g/L sodium benzoate; ····· 0.5 g/L sodium benzoate

FIGURE 9. Changes in the thermogram of liquid egg white by addition of sodium benzoate.

TABLE 2. Changes in calorimetric properties of liquid egg products by preservatives.

pH of liquid egg products and concentration of added preservative	Liquid egg white		Liquid egg yolk	Liquid whole egg	
	Peak 1	Peak 2			
Native liquid egg	T(baseline)	60.34	71.93	69.14	70.49
	T(max)	63.52	77.06	78.35	77.91
	ΔH_d	0.293	1.075	1.725	1.938
0.1 g/L K-sorbate	T(baseline)	59.88	71.27	68.23	67.93
	T(max)	63.28	76.86	77.80	77.35
	ΔH_d	0.300	0.975	1.667	1.967
0.3 g/L K-sorbate	T(baseline)	59.79	70.27	69.64	68.54
	T(max)	62.94	76.50	77.32	76.90
	ΔH_d	0.298	1.008	1.613	1.843
0.5 g/L K-sorbate	T(baseline)	59.29	70.26	66.45	67.98
	T(max)	62.57	76.03	77.16	76.74
	ΔH_d	0.295	1.016	1.725	1.912
0.1 g/L Na-benzoate	T(baseline)	60.05	71.64	69.96	70.29
	T(max)	63.52	80.96	78.34	77.89
	ΔH_d	0.282	0.950	1.822	1.924
0.3 g/L Na-benzoate	T(baseline)	59.35	70.33	69.55	70.23
	T(max)	62.74	79.97	78.15	77.83
	ΔH_d	0.288	0.960	1.750	2.063
0.5 g/L Na-benzoate	T(baseline)	59.22	71.36	69.28	66.49
	T(max)	62.51	76.34	78.01	71.92
	ΔH_d	0.287	1.037	1.675	2.023

T(baseline) – baseline denaturation temperature (°C); T(max) – peak denaturation temperature (°C); ΔH_d – change in denaturation enthalpy (J/g).

sodium benzoate content. For example, the baseline denaturation temperature of whole egg changed from 70.5°C to 66.5°C and peak denaturation temperature from around 78°C to 72°C. Potassium sorbate at the concentration of 0.5 g/L has changed denaturation temperature of liquid whole egg and liquid egg yolk by more than 2°C compared to baseline.

CONCLUSIONS

Our tests have demonstrated that preservatives used in acceptable concentrations change the protein structure and thereby calorimetric properties of liquid egg products, *i.e.* baseline denaturation temperature. The measurements have shown significant changes in calorimetric parameters of liquid egg samples by decreasing the pH value to 5.0. In addition to the reduced enthalpy due to acidification of samples, a decrease in denaturation temperature was also observed in the egg white. While the native egg white started to precipitate at 60°C, at pH 5.0 the denaturation was already started at 54.5°C. When sodium benzoate and potassium sorbate were added to liquid egg products a significant change of the tested calorimetric values was only found in the liquid egg samples containing the preservative at the concentration of 0.5 g/L. Addition of preservatives to the native liquid egg prior to treatment should be considered in technologies.

TABLE 3. 95% limits of confidence interval.

		Liquid egg white				Liquid egg yolk		Liquid whole egg	
		Peak 1	Peak 2						
pH		*	**	*	**	*	**	*	**
Native	T(baseline)	60.29	60.39	69.63	74.23	67.41	70.87	69.36	71.62
	T(max)	63.48	63.56	75.36	78.76	76.24	80.46	76.8	79.02
	ΔH_d	0.270	0.274	1.070	1.080	1.723	1.727	1.735	1.742
pH 5.5	T(baseline)	59.66	60.34	68.66	72.5	63.86	65.46	66.88	69.14
	T(max)	61.8	63.14	74.92	78.00	76.58	78.92	77	78.88
	ΔH_d	0.132	0.170	1.056	1.064	1.44	1.452	1.686	1.722
pH 5.0	T(baseline)	53.55	55.39	66.87	68.67	63.72	65.12	65.31	68.77
	T(max)	56.08	58.54	71.43	76.03	71.57	76.23	73.1	77.16
	ΔH_d	0.128	0.146	1.111	1.120	1.010	1.104	1.214	1.232
pH 4.5	T(baseline)	42.72	47.52	62.67	64.77	63.2	65.3	62.73	66.83
	T(max)	48.29	49.69	69.08	71.48	69.59	71.99	74.49	79.03
	ΔH_d	0.091	0.094	0.962	0.968	0.437	0.443	0.973	0.977
Concentration of added preservative		*	**	*	**	*	**	*	**
Native	T(baseline)	59.11	61.57	70.40	73.46	67.79	70.49	69.28	71.7
	T(max)	62.03	65.01	75.70	78.42	76.86	79.84	76.92	78.9
	ΔH_d	0.286	0.299	1.067	1.083	1.716	1.734	1.932	1.944
0.1 g/L K-sorbate	T(baseline)	57.75	62.01	69.6	72.94	66.52	69.94	66.37	69.49
	T(max)	61.34	65.22	75.04	78.68	75.92	79.68	75.73	78.97
	ΔH_d	0.291	0.308	0.967	0.982	1.657	1.678	1.870	2.064
0.3 g/L K-sorbate	T(baseline)	57.92	61.66	68.65	71.89	67.89	71.39	67.11	69.97
	T(max)	60.96	64.92	74.74	78.26	75.83	78.81	75.11	78.69
	ΔH_d	0.290	0.306	0.997	1.019	1.603	1.624	1.832	1.855
0.5 g/L K-sorbate	T(baseline)	58.62	59.96	68.57	71.95	65.62	67.28	66.95	69.01
	T(max)	61.75	63.39	73.09	78.97	75.68	78.64	75.12	78.36
	ΔH_d	0.289	0.301	1.009	1.023	1.717	1.734	1.904	1.919
0.1 g/L Na-benzoate	T(baseline)	58.49	61.61	69.58	73.70	68.2	71.72	68.58	72.0
	T(max)	61.89	65.15	78.73	83.19	76.71	79.97	76.24	79.54
	ΔH_d	0.275	0.290	0.946	0.955	1.815	1.829	1.913	1.935
0.3 g/L Na-benzoate	T(baseline)	57.69	61.01	68.97	71.69	67.81	71.29	69.25	71.21
	T(max)	61.25	64.23	78.33	81.61	76.53	79.77	76.96	78.7
	ΔH_d	0.286	0.295	0.952	0.968	1.737	1.762	1.990	2.136
0.5 g/L Na-benzoate	T(baseline)	57.35	61.09	69.44	73.28	68.05	70.51	65.78	67.20
	T(max)	60.99	64.03	74.53	78.15	76.94	79.08	71.25	72.59
	ΔH_d	0.278	0.294	1.025	1.049	1.665	1.685	1.944	2.102

T(baseline) – baseline denaturation temperature (°C); T(max) – peak denaturation temperature (°C); ΔH_d – change in denaturation enthalpy (J/g);* – lower and ** – upper limits of confidence interval (95%).

In conclusion, the calorimetric test method has proved to be useful in determinations of the effect of preservatives added to liquid whole egg as well as to liquid egg yolk and liquid egg white.

REFERENCES

1. Andr ssy E., Farkas J., Seregely Zs., Dalmadi I., Tuboly E., Lebovics V., Changes of hen eggs and their components caused by non-thermal pasteurizing treatments II. Some non-microbiological effects of gamma irradiation or hydrostatic pressure processing on liquid egg white and egg yolk. *Acta Aliment.*, 2006, 35, 305–318.
2. Carmona J.A., Cordobes F., Guerrero A., Martinez I., Portal P., Influence of pH and ionic strength on the thermal induced transitions of egg yolk dispersions. *Grasas Aceites*, 2007, 58, 289–296.
3. Chang C.M., Powrie W.D., Fennema O., Microstructure of egg yolk. *J. Food Sci.*, 1977, 42, 1193–1200.

4. Cunningham F.E., Lineweaver H., Stabilization of egg white proteins to pasteurisation temperatures above 60°C. *Food Technol.*, 1965, 19, 136–141.
5. Donovan J.W., Mapes C.J., A differential scanning calorimetric study of conversion of ovalbumin to S-ovalbumin in eggs. *J. Sci. Food Agric.*, 1976, 27, 197.
6. Donovan J.W., Mapes C.J., Davis J.G., Garibaldi J.A., A differential scanning calorimetric study of the stability of egg white to denaturation. *J. Sci. Food Agric.*, 1975, 26, 73–83.
7. Ferreira M., Hofer C., Raemy A., A calorimetric study of egg white proteins. *J. Therm. Anal.*, 1997, 48, 683–690.
8. Gliemmo K., Campos C.A., Gerschenson L.N., Effect of sweet humectants on stability and antimicrobial action of sorbate. *J. Food Sci.*, 2004, 69, 39–44.
9. Gossett P.W., Rizvi S.S.H., Baker R.C., Quantitative analysis of gelation in egg protein systems. *Food Technol.*, 1984, 38, 67–96.
10. Hammershøj M., Larsen L.B., Ipsen R.H., Qvist K.B., Effect of hen egg production and protein composition on textural properties of egg albumen gels. *J. Text. Stud.*, 2007, 32, 105–129.
11. Ichikawa T., Shimomura M., Effects of sodium chloride and other salts on the properties of diluted egg white sols and gels. *Food Sci. Technol. Int.*, 2007, 13, 173–177.
12. Landfeld A., Nesvadba P., Kyhos K., Novotna P., Pruchova J., Houska M., Sorption and thermal properties of dried egg whites. *J. Food Eng.*, 2008, 87, 59–63.
13. Lee S.Y., Kang D.H., Combined effects of heat, acetic and salt for inactivating *Escherichia coli* 0157:H7 laboratory media. *Food Contr.*, 2009, 20, 1006–1012.
14. Mohácsi-Farkas Cs., Farkas J., Mészáros L., Reichart O., Andrassy Á., Thermal denaturation of bacterial cells examined by differential scanning calorimetry. *J. Therm. Anal. Calorim.*, 1999, 57, 409–417.
15. Mizutani K., Chen Y., Yamashita H., Hirose M., Aibara S., Thermostabilization of ovotransferrin by anions for pasteurisation of liquid egg white. *Biosci. Biotechnol. Biochem.*, 2006, 70, 1839–1845.
16. Nielsen H., Hen age and fatty acid composition of egg yolk lipid. *Brit. Poultry Sci.*, 1998, 39, 53–56.
17. Ogihova H., Yatuzuka M., Horie N., Furukawa S., Yamasaki M., Synergistic effect of high hydrostatic pressure treatment and food additives on the inactivation of *Salmonella enteritidis*. *Food Contr.*, 2009, 20, 963–966.
18. Rangan C., Barceloux D.G., Food additives and sensitives. *Disease-a-month*, 2009, 55, 292–311.
19. Rossi M., Schiraldi A., Thermal denaturation and aggregation of egg proteins. *Thermochim. Acta*, 1992, 199, 115–123.
20. Torregiani A., Di Foggia M., Manco I., De Maio A., Markarian S.A., Bonora S., Effect of sulfoxides on the thermal denaturation of hens lysozyme: A calorimetric and Raman study. *J. Mol. Struct.*, 2009, 891, 115–122.
21. Zhang H., Takenaka M., Isobe S., DSC and electrophoretic studies on soymilk protein denaturation. *J. Therm. Anal. Calorim.*, 2004, 75, 719–726.

Received November 2009. Revisions received February and March 2010, and accepted June 2010.