

Pithecolobium dulce (Roxb.) Benth. Fruit Flour Intake Enhances Short-Chain Fatty Acid Production and Glucose Metabolism in BALB/c Mice

by

A.R. Hernández García, D.S. Aguilar Ávila, J.M. Viveros Paredes, R.I. López Roa, A.H. Martínez Preciado

Supplementary Materials

Table S1. Nutrient content of the standard diet Nutri-cubos brand Purina, Mexico.

Figure S1. Glucose dose-response evaluation of the positive control (inulin). (A) Oral glucose tolerance test (OGTT) at low (6.25 mg/kg), medium (25.00 mg/kg), and high (50.00 mg/kg) doses. (B) Area under the curve (AUC). Statistical analysis: one-way analysis of variance and Tukey's *post hoc* test. Significant differences: *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$. VC, vehicle control.

Table S1. Nutrient content of the standard diet Nutri-cubos brand Purina, Mexico.

Component	Content (%)
Kilocalories (kcal/g)	3.4
Carbohydrates	74.0
Protein	23.0
Fat	3.0
Total fiber	6.0
Calcium	1.0
Phosphorus	0.6
Ashes	7.0
Moisture	12.0

*Nutritional information reported by the supplier.



A**B**

Figure S1. Glucose dose-response evaluation of the positive control (inulin). (A) Oral glucose tolerance test (OGTT) at low (6.25 mg/kg), medium (25.00 mg/kg), and high (50.00 mg/kg) doses. (B) Area under the curve (AUC). Statistical analysis: one-way analysis of variance and Tukey's *post hoc* test. Significant differences: *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$. VC, vehicle control.